




PCT/CH 2004/000604



INVESTOR IN PEOPLE

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)


The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules:



Signed

Dated 13 August 2004

BEST AVAILABLE COPY

Patents Form 1/77

Patents Act 1977
(Rule 16)

THE PATENT OFFICE
DL
30 SEP 2003
RECEIVED BY FAX

1/77

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Confirmation by Mail

Cardiff Road
Newport
South Wales
NP10 8QQ

0322750.1

30 SEP 2003

1. Your reference

30090 GB

2. Patent application number
(The Patent Office will fill in this part)

0322750.1

30090 GB 30090 GB 1 30090
P01/7700 0.00-0322750.1

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Givaudan SA
Chemin de la Parfumerie 5
1214 Vernier
Switzerland

08408031001

Patents ADP Number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

Switzerland

4. Title of the Invention

ORGANIC COMPOUNDS

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Centre for Innovative Technology (Givaudan UK Ltd.)
76-80 Church Street, Staines
Middlesex TW18 4XR
United Kingdom

Patents ADP number (if you know it)

08447815001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day/month/year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application.

Number of earlier application

Date of filing
(day/month/year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:)

a) any applicant named in part 3 is not an inventor, or
b) there is an inventor who is not named as an applicant, or

c) any named applicant is a corporate body.
See note (d)

yes

Patents Form 1/77

0081303 30-Sep-03 08:40

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 9 /

Claim(s) 3 *DL*

Abstract 1 /

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77) 1 /

Request for substantive examination (Patents Form 10/77)

Any other documents
(please specify)

11. I/We request the grant of a patent on the basis of this application.

Signature

Claudia Sivet

Date 30/09/03

12. Name and daytime telephone number of person to contact in the United Kingdom

Colin Brown (office time) Tel. No: 01/784417721

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 5 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505
- b) Write your answers in capital letters using black ink or you may type them.
- c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- d) If you have answered 'Yes' Patents Form 7/77 will need to be filed
- e) Once you have filled in the form you must remember to sign and date it.
- f) For details of the fee and ways to pay please contact the Patent Office

Patents Form 1/77

DUPLICATE

1


Organic Compounds

The present invention relates to 3-isopropyl-1-methylcyclopentyl derivatives, namely (3-isopropyl-1-methylcyclopentyl)ethanol, (3-isopropyl-1-methylcyclopentyl)ethanone and (3-isopropyl-1-methylcyclopentyl)methanol and their use as fragrances. This invention relates furthermore to a method for their production and to fragrance compositions comprising them.

In the fragrance industry there is a constant demand for new compounds that enhance or improve on odour notes, or impart new odour notes.

It has now been found that certain 3-isopropyl-1-methylcyclopentyl derivatives have much sought-after floral, fruity and woody odour notes, and they are relatively simple and easy to prepare starting from naturally available (1S)-(+)- and (1R)-(-)-fenchone.

Accordingly, the present invention refers in one of its aspects to the use of a compound of formula Ia and the enantiomer (1S,3R)- thereof as fragrance



wherein

R¹ is hydrogen or methyl;

R² is hydrogen; and


R³ is hydroxyl; or

R² and R³ form together with the carbon atom to which they are attached a carbonyl group.

It has been found that the odour threshold of certain compounds of formula Ia is on an average two times lower than that of the corresponding enantiomer. Accordingly, a compound of formula I

30090 GB /29.09.03

2



enriched in its (1*R*,3*S*) enantiomer of formula Ia are preferred.

The term "enriched" is used herein to describe a compound having an enantiomeric
 5 purity greater than 1:1 in favour of the selected enantiomer. Compounds are preferred
 having a purity of about 1:3 or greater, e.g. 1:4. Particularly preferred are compounds
 having an enantiomeric purity of 1:9 or greater, such as 5:95 or 1:99.

Particularly preferred compounds of the present invention are [(1*R*,3*S*)-3-isopropyl-1-
 10 methylcyclopentyl]methanol, 1-[(1*R*,3*S*)-3-isopropyl-1-methylcyclopentyl]ethanone, and
 1-[(1*R*,3*S*)-3-isopropyl-1-methylcyclopentyl]ethanol.

The compounds according to the present invention may be used alone or in
 combination with a base material. As used herein, the "base material" includes all
 15 known odourant molecules selected from the extensive range of natural products and
 synthetic molecules currently available, such as essential oils, alcohols, aldehydes and
 ketones, ethers and acetals, esters and lactones, macrocycles and heterocycles,
 and/or in admixture with one or more ingredients or excipients conventionally used in
 conjunction with odourants in fragrance compositions, for example, carrier materials,
 20 and other auxiliary agents commonly used in the art.

The following list comprises examples of known odourant molecules, which may be
 combined with the compounds of the present invention:

- 25 – ethereal oils and extracts, e.g. tree moss absolute, basil oil, castoreum, costus root
 oil, myrtle oil, oak moss absolute, geranium oil, jasmin absolute, patchouli oil, rose oil,
 sandalwood oil, wormwood oil, lavender oil or ylang-ylang oil;
- alcohols, e.g. citronellol, EbanoTM, eugenol, farnesol, geraniol, Super MuguetTM,
 30 linalool, phenylethyl alcohol, SandaloreTM, terpineol or TimberolTM.

30090 GB /29.09.03

- aldehydes and ketones, e.g. α -amylcinnamaldehyde, Georgywood™, hydroxycitronellal, Iso E Super®, Isoraldeine®, Hedione®, maltol, Methyl cedryl ketone, methylionone or vanillin;
- 5 - ethers and acetals, e.g. Ambrox™, geranyl methyl ether, rose oxide or Spirambrene™.
- esters and lactones, e.g. benzyl acetate, Cedryl acetate, γ -decalactone, Helvetolide®, γ -undecalactone or Vetivenyl acetate.
- 10 - macrocycles, e.g. Ambrettolide, Ethylene brassylate or Exaltolide®.
- heterocycles, e.g. isobutylchinoline.
- 15 The compounds of the present invention may be used in a broad range of fragrance applications, e.g. in any field of fine and functional perfumery, such as perfumes, household products, laundry products, body care products and cosmetics. The compounds can be employed in widely varying amounts, depending upon the specific application and on the nature and quantity of other odourant ingredients. The proportion
- 20 is typically from 0.001 to 20 weight percent of the application. In one embodiment, compounds of the present invention may be employed in a fabric softener in an amount of from 0.001 to 0.05 weight percent. In another embodiment, compounds of the present invention may be used in fine perfumery in amounts of from 0.1 to 20 weight percent, more preferably between 0.1 and 5 weight percent. However, these values are
- 25 given only by way of example, since the experienced perfumer may also achieve effects or may create novel accords with lower or higher concentrations.

The compounds of the present invention may be employed into the fragrance application simply by directly mixing the fragrance composition with the fragrance

30 application, or they may, in an earlier step be entrapped with an entrapment material, for example, polymers, capsules, microcapsules and nanocapsules, liposomes, film formers, absorbents such as carbon or zeolites, cyclic oligosaccharides and mixtures thereof, or they may be chemically bonded to substrates, which are adapted to release the fragrance molecule upon application of an external stimulus such as light, enzyme,

35 or the like, and then mixed with the application.

30090 GB /29.09.03

Thus, the invention additionally provides a method of manufacturing a fragrance application, comprising the incorporation of a compound of formula I enriched in one of their enantiomers, as a fragrance ingredient, either by directly admixing the compound to the application or by admixing a fragrance composition comprising a compound of formula I enriched in one of their enantiomers, which may then be mixed to a fragrance application, using conventional techniques and methods.


As used herein, "fragrance application" means any product, such as fine perfumery, e.g. perfume and eau de toilette; household products, e.g. detergents for dishwasher, surface cleaner; laundry products, e.g. softener, bleach, detergent; body care products, e.g. shampoo, shower gel; and cosmetics, e.g. deodorant, vanishing creme, comprising an odourant. This list of products is given by way of illustration and is not to be regarded as being in any way limiting.

Compounds of formula Ia and the enantiomers thereof may be prepared by the Haller-Bauer rearrangement of (1*R*)-(-)-fenchone / (1*S*)-(+)-fenchone (1,3,3-trimethyl-2-norbornanone) followed by hydrolysis to 3-isopropyl-1-methylcyclopentanecarboxylic acid under alkali conditions, e.g. in the presence of a base such as NaOH or KOH. The resulting acid will then be reacted with methyllithium to give a compound of formula I wherein R² and R³ form together with the carbon atom to which they are attached a carbonyl group. To give further compounds of the present invention, the resulting ketone may be transformed to a tertiary alcohol either through reduction with e.g. NaBH₄, or by adding a Grignard reagent.

(3-Isopropyl-1-methylcyclopentyl)methanol may be prepared by reduction of 3-isopropyl-1-methylcyclopentanecarboxylic acid (A), which has been prepared by rearrangement of fenchone, in the presence of LAH to the corresponding alcohol, as shown in scheme 1.

5

Scheme 1:



Optically pure compounds of formula 1a and 1b and enantiomeric mixtures of a
 5 compound of formula I enriched in one of the enantiomers, i.e. a compound of formula 1a or 1b, may be synthesised, starting from optically pure fenchone or an enantiomeric mixture enriched in either (1*R*)-(-)-fenchone or (1*S*)-(+)-fenchone.

The invention is now further described with reference to the following non-limiting
 10 examples.

All end products described in the following Examples 1 to 6 are colourless oils. They
 were obtained starting from (1*R*)-(-)- and (1*S*)-(+)-fenchone that contained 8% and 2%
 respectively of the other enantiomer. The reported NMR data were measured under
 15 the following general conditions: ¹H at 400 and ¹³C at 100 MHz; in CDCl₃; chemical
 shifts (δ) in ppm downfield from TMS; coupling constants *J* in Hz.

Example 1: [(1*R*,3*S*)-3-Isopropyl-1-methylcyclopentyl]methanol

A solution of (1*R*,3*S*)-3-Isopropyl-1-methylcyclopentanecarboxylic acid (70.0 g, 0.41
 20 mol), obtained from (1*R*)-(-)-fenchone (V. Braun, J.; Jacob, A. *Chem. Ber.* 1933, 66,
 1461) in diethyl ether (100 ml) was slowly added, under nitrogen, to a suspension of
 lithium aluminium hydride (13.3 g, 0.35 mol) in the same solvent (500 ml). After heating
 at reflux during 3 h, the reaction mixture was cooled down to 10°C, 2N NaOH solution
 (70 ml) was carefully added and stirring continued for 0.5 h. The white solid was filtered
 25 off, the filtrate washed with brine (2 x 500 ml), dried (MgSO₄) and concentrated *in*
vacuo. The crude product (79.0 g) was purified by distillation using a 10 cm Vigreux
 column (0.9-1.1 mbar), 96-98°C to give [(1*R*,3*S*)-3-isopropyl-1-
 methylcyclopentyl]methanol (57.0 g, 90% yield).

30090 GB /29.09.03

¹H-NMR: δ 0.87 (d, $J = 6.7$, 3H), 0.88 (d, $J = 6.7$, 3H), 1.01 (s, 3H), 1.08 (dd, $J = 12.3$, 11.0, 1H), 1.16-1.38 (m, 3H), 1.48 (ddd, $J = 12.4$, 6.9, 0.8, 1H), 1.53-1.72 (m, 3H), 1.74-1.87 (m, 1H), 3.35 (d, $J_{AB} = 10.4$, 1H), 3.39 (d, $J_{AB} = 10.4$, 1H), ¹³C-NMR: δ 21.5 (2q),
5 25.0 (q), 30.4 (t), 33.8 (d), 35.6 (t), 41.5 (t), 43.8 (s), 46.9 (d), 72.1 (t). $[\alpha]_D^{22} -12.0$ (c 1.0, EtOH).

Odour description: floral, green, jasmine, lily-of-the-valley, fresh, clean.

10 Example 2: [(1S,3R)-3-isopropyl-1-methylcyclopentyl]methanol

Prepared according to the experimental procedure of Example 1 starting from (1S)-(+)-fenchone.

$[\alpha]_D^{22} +13.5$ (c 1.0, EtOH).

15

Odour description: floral, fruity, green, rosy, hesperidic (grapefruit).

Example 3: 1-[(1R,3S)-3-isopropyl-1-methylcyclopentyl]ethanone

A 1.6M solution of methyl lithium in diethyl ether (200 ml, 0.32 mol) was added dropwise
20 during 25 min. into a solution of (1R,3S)-3-isopropyl-1-methylcyclopentanecarboxylic acid (25.5 g, 0.15 mol) in THF (250 ml) at 0°C. After stirring at 0°C for 3 h, chlorotrimethylsilane (151 ml, 1.2 mol) was added with cooling and the reaction mixture was allowed to warm up to room temperature, poured on ice-cold water (200 ml), stirred for 0.5 h and extracted with MTBE (2 x 250 ml). The combined organic phases were
25 washed with water (200 ml), 2M NaOH (150 ml) and brine (3 x 200 ml), dried (MgSO₄) and concentrated in vacuo to give the crude [(1R,3S)-3-isopropyl-1-methylcyclopentyl]ethanone (27.6 g), a sample of which (1.5 g) was purified by bulb-to-bulb distillation (0.93 g, 68% yield).

30 ¹H-NMR: δ 0.89 (2d, $J = 6.6$, 6H), 1.19 (s, 3H), 1.24 (dq, $J = 12.4$, 9.1, 1H), 1.34-1.43 (m, 2H), 1.56-1.77 (m, 3H), 1.81-1.90 (m, 1H), 2.09 (ddd, $J = 13.1$, 9.1, 4.0, 1H), 2.15 (s, 3H). ¹³C-NMR: δ 21.3 (q), 21.4 (q), 25.0 (q), 25.3 (q), 30.2 (t), 33.3 (d), 35.6 (t), 41.0 (t), 46.6 (d), 55.3 (s), 213.0 (s). $[\alpha]_D^{22} -1.0$ (c 1.1, EtOH).

Odour description: earthy/mossy, green, woody.

Example 4: 1-[(1S,3R)-3-isopropyl-1-methylcyclopentyl]ethanone

- 5 Prepared according to the experimental procedure of Example 3 starting from (1S)-(+)-fenchone.

$[\alpha]_D^{22} +1.0$ (c 1.1, EtOH).

- 10 Odour description: floral, agrestic, fruity, green.

Example 5: 1-[(1R,3S)-3-isopropyl-1-methylcyclopentyl]ethanol

- A solution of 1-[(1R,3S)-3-isopropyl-1-methylcyclopentyl]ethanone from Example 3 (3.0 g, 18 mmol) in ethanol (8 ml) was added to a cold (ice-bath) solution of sodium borohydride (0.42 g, 10.7 mmol) in the same solvent (17 ml). After 1.5 h stirring at room temperature, the reaction mixture was poured on ice-cold 2M HCl (100 ml) and extracted with MTBE (2 x 100 ml). The combined organic phases were washed with brine (2 x 50 ml), dried (MgSO₄) and concentrated in vacuo. The crude product (2.8 g) was purified by bulb-to-bulb distillation (2.34 g, 77% yield, diastereoisomer ratio ~1:1).

- ¹H-NMR: δ 0.87 (d, J = 6.6, 3H), 0.875 (d, J = 6.6, 3H), 0.88 (2d, J = 6.6, 6H), 0.92 (s, 3H), 0.93 (s, 3H), 1.05 (t, J = 11.7, 1H), 1.12 (d, J = 6.4, 3H), 1.125 (d, J = 6.4, 3H), 1.14 (t, J = 11.8, 1H), 1.17-1.74 (m, 12H), 1.47 (2s, 2H), 1.78-1.88 (m, 2H), 3.53 (q, J = 6.3, 1H), 3.55 (q, J = 6.3, 1H). ¹³C-NMR: δ 18.5 (2q), 21.3 (2q), 21.4 (3q), 21.5 (q), 29.8 (t), 30.0 (t), 33.7 (2d), 35.8 (t), 35.9 (t), 41.9 (2t), 46.3 (2d), 46.8 (s), 46.9 (s), 75.4 (d), 75.7 (d). $[\alpha]_D^{22} -7.0$ (c 1.0, EtOH).

Odour description: floral, earthy/mossy, slightly terpineol/earthy.

Example 6: 1-[(1S,3R)-3-isopropyl-1-methylcyclopentyl]ethanol

Prepared according to the experimental procedure of Example 5 starting from (1S)-(+)-fenchone.

30090 GB /29.09.03

$[\alpha]_D^{22} +8.0$ (c 1.0, EtOH).

Odour description: hesperidic/citrus, fruity, green, fresh (grapefruit, rhubarb).

5

Example 7: Feminine Fine Fragrance

<u>Ingredient*</u>		<u>Parts per weight</u>
	Citronellol	50
10	Cyclamen aldehyde	15
	Diethyl malonate	5
	Dipropylene glycol (DPG)	149
	Florhydral	12
	Gardenol	10
15	Geraniol	50
	Hedione	25
	alpha-Hexylcinnamaldehyde	200
	Hydroxycitronellal	35
	Isocyclocitral 1% in DPG	15
20	Isojasnone	2
	Jasmal	40
	Jasmonyl	20
	Lemon oil	10
	Lilial	25
25	Linalool	65
	Linalyl acetate	50
	Methyl dianthylis	2
	Petitgrain Paraguay oil	5
	Phenethyl alcohol	65
30	Silvial	100
	<u>[(1R,3S)-3-Isopropyl-1-methylcyclopentyl]methanol</u>	<u>50</u>
		Total 1000

30090 GB /29.09.03

* for chemical names see Flavor and Fragrance Materials — 2003, Allured Publishing Corp. Carol Stream Ill., U.S.A..

The presence of 5% of [(1*R*,3*S*)-3-Isopropyl-1-methylcyclopentyl]methanol confers to this formula a *creamy, lily-of-the-valley* aspect.

Example 8: Floral Composition for Soap

10	<u>Ingredient</u>	<u>Parts per weight</u>
	Agrumex	100
	Benzophenone	60
	Benzyl acetate	55
	Bergamot base	80
15	4- <i>t</i> -Butylcyclohexyl acetate	150
	Diphenyl oxide	20
	Dipropylene glycol (DPG)	78
	Ebanol	20
	Hydroxycitronellal	200
20	Jasmine base	80
	Methyl Phenylacetate	2
	Nerol	20
	Phenylpropyl alcohol	40
	Rose base	100
25	Rhodinol	65
	Sandela	30
	Silvial	100
	<u>[(1<i>R</i>,3<i>S</i>)-3-Isopropyl-1-methylcyclopentyl]methanol</u>	<u>50</u>
	Total	1250


* for chemical names see Flavor and Fragrance Materials — 2003, Allured Publishing Corp. Carol Stream Ill., U.S.A..

[(1*R*,3*S*)-3-Isopropyl-1-methylcyclopentyl]methanol makes this *lily-of-the-valley* fragrance velvety and rich.

30090 GB /29.09.03

Claims

1. The use of a compound of formula Ia and the enantiomer thereof as fragrance,



wherein

R¹ is hydrogen or methyl;


R² is hydrogen; and

R³ is hydroxyl; or


R² and R³ form together with the carbon atom to which they are attached a carbonyl group.

2. The use of a compound according to claim 1 selected from: [(1R,3S)-3-isopropyl-1-methylcyclopentyl]methanol, [(1S,3R)-3-isopropyl-1-methylcyclopentyl]methanol, 1-[(1R,3S)-3-isopropyl-1-methylcyclopentyl]ethanone, 1-[(1S,3R)-3-isopropyl-1-methylcyclopentyl]ethanone, 1-[(1R,3S)-3-isopropyl-1-methylcyclopentyl]ethanol and 1-[(1S,3R)-3-isopropyl-1-methylcyclopentyl]ethanol.

3. The use of a compound of formula I




enriched in the enantiomer having the formula Ia



wherein R¹, R² and R³ have the same meaning as given in claim 1.


11

4. The use of a compound of formula I



I

enriched in the enantiomer having the formula Ib




Ib

(1S,3R)-

wherein R¹, R² and R³ have the same meaning as given in claim 1.

5. The use of a compound as defined in one of the preceding claims in fragrance applications.
6. A fragrance application comprising a compound as defined in any of the preceding claims 1 - 4.
7. A fragrance application according to claim 6 wherein the fragrance application is a perfume, household product, laundry product, body care product or cosmetic products.
8. A method of manufacturing a fragrance application, comprising the step of incorporating a compound of formula Ia or its enantiomer as defined in claim 1, 2, 3 and 4.
9. A compound of formula Ia



Ia

(1R,3S)-

30090 GB/29.09.03

12

wherein


R^1 is hydrogen or methyl;

R^2 is hydrogen; and

R^3 is hydroxyl; or

R^2 and R^3 form together with the carbon atom to which they are attached a carbonyl group.

10. A compound of formula Ib



Ib

(1S,3R)

wherein

R^1 is hydrogen or methyl;

R^2 is hydrogen; and

R^3 is hydroxyl; or

R^2 and R^3 form together with the carbon atom to which they are attached a carbonyl group.

Abstract

This invention relates to 3-isopropyl-1-methylcyclopentyl derivatives and their use in fragrance applications.

PCT/CH2004/000604



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.